



VANUATU AgroMet Bulletin


Vanuatu Meteorology & Geo-Hazards Department
Department of Agriculture & Rural Development



November 2021




Vol.1, Issue 9

CURRENT ENSO STATUS



ENSO outlook has been raised to La Niña ALERT. This means there is a 70% chance for La Niña to happen in the coming months. La Niña typically brings above normal rainfall over Vanuatu, with impacts such as flooding, river overflow, with possible landslide/mudslide over parts of the country.


Temperature and Rainfall Outlook for December 2021:



Summary:


- Vanuatu is expecting warmer than average day and night time temperature in December.
- Above normal rainfall is expected over the country in November with 50% to 70% chances for these forecast to happen.

Drought Status for November 2021:



- All Vanuatu regions has been and will be expecting extremely higher than average (Alert 3 wet) rainfall in this month of November.
- Forecasts also show high chances (Alert 3 Wet) of above average rainfall is very likely for the whole country in the next three months, December to February 2022.

Outlook for Dec to Feb 2022:



Top Highest and Lowest Tides for November to December 2021: Luganville & Port Vila Harbor

Luganville Harbour					
Lowest Tide	Date	Time (VUT)	Highest Tide	Date	Time (VUT)
0.19m	7 Nov	00:08am	1.91m	5 Nov	16:44pm
0.11m	6 Dec	00:00am	1.94m	4 Dec	16:22pm

Port Vila Harbour					
Lowest Tide	Date	Time (VUT)	Highest Tide	Date	Time (VUT)
0.16m	7 Nov	00:59am	1.63m	6 Nov	17:36pm
0.09m	6 Dec	00:47am	1.65m	4 Dec	17:08pm

Moon Phases for November to December 2021

New Moon	First Quarter	Full Moon	Last Quarter
5 November	11 November	19 November	27 November
4 December	11 December	19 December	27 December

Climate Smart Recommendations for crop cultivation

Crop (variety) Selection

Vegetables	<input type="checkbox"/> Depending on market and demand, and availability of seeds and seedlings. Select varieties that can perform well in Hot and wet season (November to April) e.g. Corn, cucumber, capsicum
Island Taro	<input type="checkbox"/> Select Varieties with higher demand and market and available planting material (Available varieties: Sakius, Tarapatan, Alkat) and other recommended varieties from your site, village or island.
Manioc	<input type="checkbox"/> Consider growing varieties depending on target market and demand, and available planting materials. Grow dwarf varieties due to cyclone season
Yam	<input type="checkbox"/> Depending on target market and demand, and available planting material. Select disease resistant varieties (anthracnose) for planting during this season.
Kumala	<input type="checkbox"/> Depending on target market and demand, and available planting material (Available varieties: Baby kumala, PNG, bankis, Epule 1, Fanafo, salili and kumala hybrids)

Planting

Vegetables	<input type="checkbox"/> Select healthy seeds and seedlings before planting. Avoid transplanting of seedlings during sunny mid days or during heavy rain. Consider weekly/daily weather updates through media (FB, radio etc.) to sow seeds and transplant vegetables. Traditional weather forecasting are also advice to use
Island Taro	<input type="checkbox"/> Select good type of planting material from healthy plants and free from pest and diseases, good size to produce good corm shape, large suckers or headset have rapid early growth and higher survival rate.
Manioc	<input type="checkbox"/> Select healthy stalks which are free from pest and diseases. It is best to collect stalks just after harvested and not sprouted, because sprouting before planting will weaken the ability of stalks to grow after the stalk is cut and planted. It is best to collect the middle part of the stalk as planting material.
Yam	<input type="checkbox"/> Select tubers with minimum rot; Trim any rot from tubers before cutting it up into planting pieces. Yam is not a shade tolerant plant, it is a light loving and shade sensitive plant thus require sites that are well exposed to sunlight; Staking should be low (1m height).
Kumala	<input type="checkbox"/> It is best to select cuttings from the tip of the vine about 30cm to 40cm length and which are free from Pest and diseases; collect cuttings from young plants which are 2 to 3 months of age. Avoid planting materials with leaf scab disease and also do not collect planting materials from plants with little leaf disease. Kumala are also light loving crop and prefer sites that are expose to good sunlight.

Farm Management (nutrient/pest/weed)

Vegetables	<input type="checkbox"/> Monitor/control insect pest and diseases attack during the growth stage of the crops; Plant repellent crops/plants to reduce pest and disease attack; Remove infected plants and apply intercropping; Apply weeding as needed and it can be done 2-3 weeks after planting; apply drainage to reduce flooding.
Island Taro	<input type="checkbox"/> Monitor disease symptoms and insect pests (e.g. Taro beetle, caterpillar and aphids infestation on leaves) during every farm visit; apply weeding every 2-3 weeks for up to 4 months; Apply mulching when needed using organic materials (vetiver grass) to help reduce weed growth and add organic matter to the soil; Apply intercropping system to reduce Pest and disease attack to plants, apply agroforestry system.
Manioc	<input type="checkbox"/> Apply mulching and composting when needed during the early growth stage of the crop; Monitor weed growth and weed as needed, first weeding can be done after 3-4 weeks after planting; Control major Pest such as Rat after tuber formation; Remove infected plants and apply intercropping system
Yam	<input type="checkbox"/> Weeding can be applied at least 2 or 3 times during the growth cycle, depending on the weeds present in the field. Monitor leaf eating beetles during the early growth stage of yam and apply cultural practices as needed to control the beetles. Monitor vine growth and adjust to staking accordingly
Kumala	<input type="checkbox"/> Monitor weed growth and weed as needed; Weeding will usually be every 2 weeks for up to 5-6 weeks by when the runners have covered the space between rows. Mounding should be done 2 weeks and 1 month after planting to reduce damage on stem and tubers by the weevil and rat.

CONTACTEM MIFALA:

• Department of Agriculture & Rural Development
PMB 9040
Telephone: (678) 33550. Email: Pleo@vanuatu.gov.vu

• Vanuatu Meteorological & Geo-Hazards Department
PMB 9054
Tel: (678) 24686. Email: climate@meteo.gov.vu

Supported by:

